

Orthogonal polynomials of several discrete variables and the 3nj-Wigner symbols: applications to spin networks

M Lorente

Departamento de Física, Universidad de Oviedo, 33007 Oviedo, Spain

Abstract. The use of orthogonal polynomials for integral models on the lattice is applied to the 3nj-symbols that appear in the coupling of several angular momenta. These symbols are connected to the Ponzano-Regge method to solve the Einstein equations on a discrete Riemannian manifold.

1. Classical orthogonal polynomials of one discrete variable

These polynomials satisfy a difference equation of hypergeometric type such that the difference derivatives of the some polynomial satisfy a similar equation. The discrete variable can be consider of two types.

a) *On homogeneous lattice:* $x = 0, 1, 2, \dots$

The corresponding polynomials satisfy a difference equation

$$\sigma(x)\Delta\nabla p_n(x) + \tau(x)\Delta p_n(x) + \lambda_n p_n(x) = 0$$

with $\Delta f(x) = f(x+1) - f(x)$, $\nabla f(x) = f(x) - f(x-1)$, $\sigma(x)$ and $\tau(x)$ are functions of second and first order respectively;

an orthogonality relation

$$\sum_{x=a}^{b-1} p_n(x) p_m(x) \rho(x) = d_n^2 \delta_{mn}$$

with $\rho(x)$ a weight function and d_n a normalization constant. To these polynomials correspond the Meixner, Kravchuk, Charlier and Hahn polynomials [1]

b) *On non homogeneous lattice:* $x = x(s)$, $s = 0, 1, 2, \dots$ the polynomials satisfy a difference equation

$$\sigma[x(s)] \frac{\Delta}{\Delta x \left(s - \frac{1}{2}\right)} \frac{\nabla y_n(x)}{\nabla x(s)} + \frac{1}{2} \tau[x(s)] \left\{ \frac{\Delta y_n(x)}{\Delta x(s)} + \frac{\nabla y_n(x)}{\nabla x(s)} \right\} + \lambda_n y_n(x) = 0;$$

an orthogonality relation

$$\sum_{s=a}^{b-1} y_n(s) y_m(s) \rho_n(x) \Delta x \left(s - \frac{1}{2}\right) = d_n^2 \delta_{nm}.$$

The corresponding polynomials are classified according to the lattice function:

For $x(s) = s(s+1)$, we have the Racach and dual Hahn polynomials

For $x(s) = q^s$ or $\frac{q^s - q^{-s}}{2}$, we have the q -Kravchuk, q -Meixner, q -Charlier and q -Hahn polynomials.

For $x(s) = \frac{q^s + q^{-s}}{2}$ or $\frac{q^{is} + q^{-is}}{2}$, we have the q -Racach and q -dual Hahn polynomials [2].

2. Generalized Clebsch-Gordon coefficients and generalized $3nj$ -Wigner symbols

If two angular momentum operators are coupled to give a total angular momentum $J = J_1 + J_2$ the new basis can be expressed in terms of the old ones

$$|j_1 j_2 j m\rangle = \sum_{m_1+m_2=m} \langle j_1 j_2 m_1 m_2 | j_1 j_2 j m\rangle |j_1 j_2 m_1 m_2\rangle.$$

The symmetry properties of the Clebsch-Gordon coefficients in this expansion are more patent if one substitutes them by the Wigner symbols

$$\langle j_1 j_2 j m | j_1 j_2 m_1 m_2\rangle = (-1)^{j_1-j_2+j-m} \sqrt{2j+1} \begin{pmatrix} j_1 & j_2 & j \\ m_1 & m_2 & -m \end{pmatrix}$$

Similarly if we couple three angular momentum operator we obtain a new basis in terms of the old ones:

$$|j_1 j_2 j_3 j_1 j_2 j m\rangle = \sum \langle j_1 j_2 j_3 m_1 m_2 m_3 | j_1 j_2 j_3 j_1 j_2 j m\rangle |j_1 j_2 j_3 m_1 m_2 m_3\rangle$$

for the coupling $(J_1 + J_2) + J_3 = J$,

$$|j_1 j_2 j_3 j_2 j_3 j m\rangle = \sum \langle j_1 j_2 j_3 m_1 m_2 m_3 | j_1 j_2 j_3 j_2 j_3 j m\rangle |j_1 j_2 j_3 m_1 m_2 m_3\rangle$$

for the coupling $J_1 + (J_2 + J_3) = J$.

Both bases are related by some matrix $U(j_{12}, j_{23})$ that can be written in terms of generalized $6j$ -Wigner symbol

$$U(j_{12}, j_{23}) = (-1)^{j_1+j_2+j_3+j} \sqrt{(2j_{12}+1)(2j_{23}+1)} \begin{Bmatrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{Bmatrix}$$

In similar fashion can be written the generalized Clebsch-Gordon coefficients and generalized $3nj$ -Wigner symbols [3]. The algebraic properties of these symbols can be represented by geometrical graphs [3].

3. $3nj$ -symbols as orthogonal polynomials of several discrete variable

The $6j$ -symbols are proportional to the Racah polynomials through the following relation [2]

$$(-1)^{j_1+j_2+j_{23}} \sqrt{(2j_{12}+1)(2j_{23}+1)} \begin{Bmatrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{Bmatrix} = \frac{\sqrt{\rho(x)}}{d_n} u_n^{(\alpha,\beta)}(x, a, b)$$

with

$$\begin{aligned} x(s) &= s(s+1), \quad s = j_{23} \\ a &= j_3 - j_2, \quad b = j + j_3 + 1, \quad n = j_{12} - j_1 + j_2 - j \\ \alpha &= j_1 - j_2 - j_3 + j, \quad \beta = j_1 - j_2 + j_3 - j \end{aligned}$$

Using the asymptotic limit of the Racah polynomials and the connections between the Jacobi polynomials and the Wigner little functions one can prove the following approximation of the $6j$ -symbols when $j_1 \sim j_2 \sim j_3 \sim j \gg j_{12}$

$$\begin{Bmatrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{Bmatrix} \simeq \frac{(-1)^{j_2+j_3+j_{23}}}{\sqrt{j_1+j_2+1}\sqrt{j_3+j+1}} d_{j_1-j_2, j_3-j}^{j_{12}}(\vartheta) \quad (1)$$

with

$$\cos \theta = \frac{(2j_{23} + 1)^2 - (j_1 + j_2 + 1)^2 - (j_3 + j + 1)^2}{2(j_1 + j_2 + 1)(j_3 + j + 1)}$$

The $3nj$ -symbols of the first and second kind can be written in terms of $6j$ -symbols, and therefore in terms of product of Racah polynomials, giving rise to orthogonal polynomials of several discrete variables. To illustrate this take, f.i., the $12j$ -symbol of the second kind as a combination of $6j$ -symbols.

$$\left\{ \begin{matrix} j_1 & j_2 & j_3 & j_4 \\ l_1 & l_2 & l_3 & l_4 \\ k_1 & k_2 & k_3 & k_4 \end{matrix} \right\} = \sum_x (2x + 1) (-1)^{R_n + 4x} \left\{ \begin{matrix} j_1 & k_1 & x \\ k_2 & j_2 & l_1 \end{matrix} \right\} \left\{ \begin{matrix} j_2 & k_2 & x \\ k_3 & j_3 & l_2 \end{matrix} \right\} \left\{ \begin{matrix} j_3 & k_3 & x \\ k_4 & j_4 & l_3 \end{matrix} \right\} \left\{ \begin{matrix} j_4 & k_4 & x \\ k_1 & j_1 & l_4 \end{matrix} \right\}$$

Here $R_n = \sum_{i=1}^4 (j_i + l_i + k_i)$. Substituting each $6j$ -symbol for the corresponding Racah polynomial we obtain:

$$\left\{ \begin{matrix} j_1 & j_2 & j_3 & j_4 \\ l_1 & l_2 & l_3 & l_4 \\ k_1 & k_2 & k_3 & k_4 \end{matrix} \right\} = \sum_x \frac{1}{2x + 1} \prod_{i=1}^4 \frac{\sqrt{\rho(l_i)}}{d_{n_i}} u_{n_i}^{(\alpha_i, \beta_i)}(l_i) \equiv p_n(l_1 \ l_2 \ l_3 \ l_4)$$

which is a polynomial of four discrete variables.

For the asymptotic limit we find

$$\left\{ \begin{matrix} j_1 & j_2 & j_3 & j_4 \\ l_1 & l_2 & l_3 & l_4 \\ k_1 & k_2 & k_3 & k_4 \end{matrix} \right\} \approx \sum_x (2x + 1) \prod_{i=1}^4 \frac{1}{j_i + k_i + 1} d_{j_i - k_i, j_{i+1} - k_{i+1} \pmod{4}}^x(\vartheta_i)$$

These formulas can be easily generalized to any $3nj$ -symbols of first and second kind.

4. Application to spin networks and to Ponzano-Regge integral action

Penrose has proposed a model for the space and time in which the underlying structure is given by a set of interactions between elementary units that satisfy the coupling of angular momentum operators, called spin networks [4]. One particular case of these networks can be described by the graphs of $3nj$ -symbols. From different point of view Regge has proposed a method to calculate Einstein action by the approximation of curved riemannian manifold by a polyedron built up of triangles. Later Ponzano and Regge applied the properties of $6j$ -symbols to calculate the sum action over this triangulation [5]

Let M be a riemannian manifold that is approximated by a polyedron with boundary D and it is decomposed into p tetrahedra T_k represented by $6j$ -symbols.

The polyedron give rise to triangular faces f , represented by $3j$ -symbols, and to q internal edges x_i , as well as to external ones l_i with respect to the boundary D .

Ponzano and Regge define the sum

$$S = \sum_{x_i} \prod_{k=1}^p T_k (-1)^\varphi \prod_{i=1}^q (2x_i + 1) \quad (2)$$

When $l_i \rightarrow \infty$, $\hbar \rightarrow 0$, $\hbar l_i \rightarrow \text{finite}$ we recovered the continuous manifold. In order to compute the $6j$ -symbols in the classical limit, we use the asymptotic formula [2]

$$d_{mm'}^j(\theta) \approx (-1)^{m-m'} \sqrt{\frac{2}{\pi(j-m)}} \left(\frac{2j+m-m'+1}{2j-m+m'+1} \right)^{\frac{m+m'}{2}} \frac{\cos \left[\left(j + \frac{1}{2} \right) \theta - \left(m - m' + \frac{1}{2} \right) \frac{\pi}{2} \right]}{\sqrt{\sin \theta}}$$

at $m \sim m' \sim 1$, $j \gg 1$. Substituting this expression in (1) with $m = j_1 - j_2$, $m' = j_4 - j_5$, $j = j_6$ and taking the edges of the tetrahedra $j_1 + \frac{1}{2}, \dots, j_6 + \frac{1}{2}$, very large except j_6 , we have

$$\begin{aligned} \begin{Bmatrix} j_1 & j_2 & j_3 \\ j_4 & j_5 & j_6 \end{Bmatrix} &\approx \frac{1}{\sqrt{12\pi V}} \cos \left\{ \left(j_6 + \frac{1}{2} \right) \theta - \left(j_1 + \frac{1}{2} \right) \frac{\pi}{2} + \right. \\ &\quad \left. + \left(j_2 + \frac{1}{2} \right) \frac{\pi}{2} - \left(j_4 + \frac{1}{2} \right) \frac{\pi}{2} + \left(j_5 + \frac{1}{2} \right) \frac{\pi}{2} + \frac{\pi}{4} \right\} = \\ &= \frac{1}{\sqrt{12\pi V}} \cos \left\{ \sum_{i=1}^6 \left(j_i + \frac{1}{2} \right) \theta_i + \frac{\pi}{4} \right\} \end{aligned} \quad (3)$$

where θ_i is the dihedral angle for the edge j_i and $V = \frac{1}{6} \left(j_1 + \frac{1}{2} \right) \left(j_4 + \frac{1}{2} \right) \left(j_6 + \frac{1}{2} \right) \sin \theta$. Note the formula (3) has been proved rigorously by Roberts [5]. Introducing formula (3) in formula (2), Ponzano and Regge proved that it leads in the continuous limit to the integral action of the general relativity.

This work has been partially supported by M.I.C. (grant FM2000-0357) Spain

References

- [1] M. Lorente, "Raising and lowering operators, factorization and difference operators of hypergeometric type", *J.Phys.A.Math.Gen* 34 (2001) 569-588.
- [2] Nikiforov, A.F. et al., "Classical orthogonal polynomials of a discrete variable", Springer 1991.
- [3] Yutsis, A.P. et al., "Mathematical apparatus of the theory of angular momentum", Israel program for scientific translations, Jerusalem 1962.
- [4] Penrose, R., "Angular momentum: an approach to combinatorial space-time" in Quantum theory and Beyond" (T. Bastin ed.) Cambridge 1970.
- [5] Regge T., Williams R., "Discrete structures in gravity", *J. Math Phys.* 41 (2000) 3964-3984.